



Air Quality Monthly Report

December, 2016



Department of Environment
Ministry of Environment, Forest and Climate Change
Bangladesh

Content

1.	Introduction.....	1
2.	Monitoring Network	2
3.	Monthly Air Quality	4
4.	Summary and conclusion.....	5
5.	ANNEX.....	9

1. Introduction

Air quality management plans based on knowledge of sources, appropriate air quality standards, accurate air quality data, and effective incentives; and enforcement policies is therefore needed to be adopted.

At this backdrop, real-time measurements of ambient level pollutants were made at 8 major cities (Namely, Dhaka, Narayanganj, Gazipur, Chittagong, Rajshahi, Khulna, Barisal and Sylhet) of Bangladesh. The data generated will be used to define the nature and severity of pollution in the cities; identify pollution trends in the country; and develop air models and emission inventories.

The program encompasses operation of the sampling and monitoring network, and quality assurance activities to ensure the quality of the data collected and disseminated by the CASE project.

CASE project monitors the criteria pollutants such as carbon monoxide, nitrogen dioxide, ozone, sulfur dioxide, PM10 and PM2.5. Monitoring is performed to demonstrate attainment or non-attainment of national ambient air quality standards to assess the trends of air pollution levels.

The main purpose of this report is to present, analyze and make available of these data to the general public, stakeholders, researchers and policy makers to develop effective air pollution abatement strategies. This report summarizes the air quality data collected at the different CAMS in operation under the Department of Environment (DoE) air quality monitoring network.

The basis for discussion of air quality has been the data collected from the Air Quality monitoring Network stations under DoE. The data have been quality controlled and the air pollution levels have been compared to the Bangladesh Ambient Air Quality Standard as adopted in 2005. Table 1 represents the current and approved air quality standards for Bangladesh.

Table 1: National Ambient Air Quality Standards for Bangladesh

Pollutant	Objective	Average
CO	10 mg/m ³ (9 ppm)	8 hours(a)
	40 mg/m ³ (35 ppm)	1 hour(a)
Pb	0.5 µg/m ³	Annual
NO _x	100 µg/m ³ (0.053 ppm)	Annual
PM10	50 µg/m ³	Annual (b)
	150 µg/m ³	24 hours (c)
PM2.5	15 µg/m ³	Annual
	65 µg/m ³	24 hours
O ₃	235 µg/m ³ (0.12 ppm)	1 hour (d)
	157 µg/m ³ (0.08 ppm)	8 hours
SO ₂	80 µg/m ³ (0.03 ppm)	Annual
	365 µg/m ³ (0.14 ppm)	24 hours (a)

Notes:

- (a) Not to be exceeded more than once per year
- (b) The objective is attained when the annual arithmetic mean is less than or equal to 50 µg/m³
- (c) The objective is attained when the expected number of days per calendar year with a 24-hour average of 150 µg/m³ is equal to or less than 1
- (d) The objective is attained when the expected number of days per calendar year with the maximum hourly average of 0.12 ppm is equal to or less than 1 (Source: AQMP, DOE).

The air quality index (AQI) is a number for reporting the state of air quality in an area on daily basis. In Bangladesh the AQI is based on 5 criteria pollutants; Particulate Matter (PM₁₀ and PM_{2.5}), NO₂, CO, SO₂ and Ozone (O₃). AQI tells how clean or polluted the air is, and what associated health effects might be a concern for public. The AQI focuses on health effects that one might experience within a few hours or days after breathing polluted air. The AQI is a single number from 0 to 500 calculated on the basis of pollutant concentration measured in an area. Higher the AQI, higher is the air pollution level and thus indicates greater public health concern.

An AQI value of 100 generally corresponds to the national air quality standard for the pollutant, which is the level that set by the mandated Environment Protection Agency (e.g., for Bangladesh Department of Environment) to protect public health. AQI values below 100 are generally thought of as satisfactory. When AQI values are above 100, air quality is considered to be unhealthy-at first for certain sensitive groups of people, then for everyone as AQI values get higher. The AQI standard for Bangladesh is given in Table 2.

Table 2: Air Quality Index (AQI) for Bangladesh

Air Quality Index (AQI) range	Category	Color
0-50	Good	Green
51-100	Moderate	Yellow Green
101-150	Caution	Yellow
151-200	Unhealthy	Orange
201-300	Very Unhealthy	Red
301-500	Extremely Unhealthy	Purple

2. Monitoring Network

The main objective of the Bangladesh AQM network is to provide reliable information to the authorities and to the public about the air quality in most populous cities of Bangladesh.

As a part of the air quality monitoring strategy, several objectives can be achieved, including:

- Establish source/receptor relationships;
- Identify which are the pollutants of concern and their current status;
- Show how widespread air pollution problems are and indicate the general extent of the public exposure;
- Provide benchmarks against which trends in overall air quality can be compared and devise performance indicators for assessing the impact of an air quality management plan or strategy;
- Provide a data base for evaluation of effects; of urban, land use management, and transportation planning; of development and evaluation of abatement strategies; and of development and validation of atmospheric processes and models.

Another objective in the monitoring and management programmes is to provide input data for modeling. These data will serve as a background for performing air quality planning and abatement studies. Model results may also serve as input to other studies such as health related investigations and exposure assessments.

The ambient air quality monitoring network Bangladesh consists of eleven (11) fixed Continuous Air Monitoring Stations (CAMS). The locations of the 11 CAMS are shown in Figure 1. Brief description of the monitoring stations and the list of measured parameters recorded at each station are provided in Table 3.

Table 3: Description of Monitoring Network:

City	ID	Location	Lat/Lon	Monitoring capacity
Dhaka	CAMS-1	Sangshad Bhaban, Sher-e-Bangla Nagar	23.76N 90.39E	PM10, PM2.5, CO, SO2, NOX, O3, and HC concentrations with meteorological parameters.
	CAMS-2	Farmgate	23.76N 90.39E	PM10, PM2.5, CO, SO2, NOX, O3, and HC with meteorological parameters.
	CAMS-3	Darus-Salam	23.78N 90.36E	PM10, PM2.5, CO, SO2, NOX and O3 with meteorological parameters.
Gazipur	CAMS-4	Gazipur	23.99N 90.42E	PM10, PM2.5, CO, SO2, NOX and O3 with meteorological parameters.
Narayangonj	CAMS-5	Narayangonj	23.63N 90.51E	PM10, PM2.5, CO, SO2, NOX and O3 with meteorological parameters.
Chittagong	CAMS-6	TV station, Khulshi	22.36N 91.80E	PM10, PM2.5, CO, SO2, NOX, O3, and HC with meteorological parameters.
	CAMS-7	Agrabad	22.32N 91.81E	PM10, PM2.5, CO, SO2, NOX and O3 with meteorological parameters.
Sylhet	CAMS-8	Red Crecent Campus	24.89N 91.87E	PM10, PM2.5, CO, SO2, NOX and O3 with meteorological parameters.
Khulna	CAMS-9	Baira	22.48N 89.53E	PM10, PM2.5, CO, SO2, NOX, O3, and HC with meteorological parameters
Rajshahi	CAMS-10	Sopura	24.38N 88.61E	PM10, PM2.5, CO, SO2, NOX, O3, and HC with meteorological parameters.
Barisal	CAMS-11	DFO office campus	22.71N 90.36E	PM10, PM2.5, CO, SO2, NOX and O3 with meteorological parameters.




Figure 1: CAMS Location in Bangladesh

Monitoring data from network stations are transferred to a central data centre at the Department of Environment office in Dhaka and simultaneously transferred to Air Quality Management System based on NILU AIRQus system established under BAPMAN project. The data are stored in AIRQus database for quality check, control, evaluation, validation, statistical analysis. Quality controlled data are then stored in the final database for further analysis, reporting, presentations and future use.

3. Monthly Air Quality

The data presented in this report are based on monitoring of air quality parameters during December, 2016 at 11 CAMS operated under CASE-DoE monitoring network. Table-4 summarizes the basic statistics of the data along with the data capture rate and the number of days for which specific pollutant exceeded the Bangladesh National Ambient Air Quality Standard (BNAQS). Since NO_x have only annual standard, so for this pollutant daily 24-hours average concentration levels were compared with the annual average. During data quality control some data, which are outliers (beyond 3rd and 97th percentile) and inconsistent data, were flagged as invalid and those were not included in the analysis. Time series plots based on the data generated in the CAMS are also given in Annexes.

Data availability (valid data) from those analyzers was functional found to be over 80% except few parameters in different CAMS in operation. During the reporting month several analyzers measuring gaseous pollutants (especially SO₂) were not operational due to routine preventive/corrective maintenance. Data from Sangsad Bhaban CAMS, Khulna CAMS and TV station, Chittagong CAMS not included in the report due to nonfunctional of Data logger computer.

Inspection of the available data shows that there were few occurrences of non-compliance for PM₁₀ & PM_{2.5} levels at the monitoring stations during the month of December, 2016. It is observed that the 24 hr average concentration level of PM_{2.5} exceed BNAQS 29-30 days at Gazipur, Barishal and Agrabad, Chittagong CAMS, 21-23 days at Darussalam, Rajshahi

and Narayonganj CAMS, 11 days at Sylhet CAMS respectively. For PM₁₀ non-attainment with respect to BNAAQS occurred for 28-29 days in Narayonganj and Gazipur CAMS, 21-24 Rajshahi and Darussalam CAMS, Agrabad Chittagong CAMS, 13-14 days in Barisal CAMS & Sylhet CAMS at during the reporting month. PM_{2.5} and PM₁₀ results are not reported in the month for TV Station Chittagong, Sangsad Bhaban, BARC, Farmgate and Khulna CAMS due to malfunction of PM Monitor and Data logger computer. The monthly average concentration level of PM_{2.5} and PM₁₀ measured at different CAMS were found 86.5-206 µg/m³ and 160-345 µg/m³ respectively during the month of December, 2016. That concentration level of those was found 49-119 µg/m³ and 100-246 µg/m³ respectively during the month of November, 2016. From the time series plot of both PM₁₀ and PM_{2.5}, it is seen there are most of the episodes of PM concentrations is high. 24-hours average PM levels in all cities monitored are found higher compared to the previous month because prevailing dry season and lower wind speed. It is also observed that all the gaseous pollutants except NO_x and CO measured at different CAMS did not exceed the BNAAQS during the month of December, 2016. NO_x concentrations exceeded the BNAAQS 17 days in Darussalam CAMS, 02-03 days in Agrabad Chittagong and Sylhet CAMS in this month. It is observed that 8 hr average CO concentration exceeded BNAAQS 19 hours at the reporting month at BARC CAMS located near main road.

In general PM pollution levels in the cities monitored during the reporting month found higher with few exceptions. Usually in the dry seasons the pollution level reached highest peak and gradually decreases during wet season, which is reflected in the data monitored in all CAMS during the month of December, 2016. It is observed that average wind speed and precipitation compared to previous month is decreased, which decreases the rate of dispersion of the pollutants and this might be a reason for observed higher PM concentration

Daily air quality index (AQI) values were calculated based on the available air quality data (valid data) from different CAMS and summary of the AQI by categories (Table:2) are presented in annex Figure 5. Summary data shows majority of the days AQI values were in very unhealthy categories or extremely unhealthy categories and few caution, moderate and unhealthy categories.

4. Summary and conclusion

Data obtained from CAMS operated under DoE air quality monitoring network during December, 2016 have been analyzed and reported. Data availability was 65-80% for all the criteria pollutant monitored at different CAMS with few exceptions. Air quality data for some pollutants were not reported because either the analyzer was not functional or the data capture rate was too low. From the analysis of the data following conclusion can be drawn:

- Although PM₁₀ and PM_{2.5} are the most critical pollutants but 24-hour average for both PM₁₀ and PM_{2.5} concentrations during reporting month were found higher. It is observed that the average concentration level of PM_{2.5} and PM₁₀ measured at different CAMS were 86.5-206 µg/m³ and 160-345 µg/m³ respectively during the month of December, 2016.
- The gaseous pollutants except NO_x measured at different CAMS did not exceed limit values of the BNAAQS. NO_x concentrations exceeded the BNAAQS 17 days in Darussalam CAMS, 02-03 days in Agrabad Chittagong and Sylhet CAMS in this month. 8 hr average CO concentration exceeded BNAAQS 19 hours at BARC CAMS.
- Due to decreasing average wind speed and precipitation during December, 2016, dispersion and wash out of pollutants decreases and thus the pollution concentration level increases.
- Monthly summary of calculated AQI values based on data from different CAMS showed that during this month most of the day's air quality was either Very Unhealthy or extremely Unhealthy categories or few caution, moderate & unhealthy categories

as well. In all cases most frequent responsible pollutant was PM2.5. In absence of PM2.5 sometimes found responsible pollutant PM10.

During the reporting month number of analyzer especially SO2 of new CAMS did not produce good data and availability of data was not satisfactory and they are under maintenance process.

Table 4: Summary Air Quality and Meteorological data measured during December, 2016 at different CAMS operated under DoE

Parameter	unit	NAAQS	Summary	CAMS-1 (S-Bhaban)	CAMS-2 (BARC) ^a	CAMS-3 (D-salam)	CAMS-4 (Gazipur)	CAMS-5 (Narayong anj)	CAMS-6 TV-St (Chittagong) ^a	CAMS-7 Agrabad-(Chittagong)	CAMS-8 (Sylhet)	CAMS-9 (Khulna) ^a	CAMS-10 (Rajshahi) ^a	CAMS-11 (Barisal)
SO ₂ -24 hr	ppb	140	Average	DNA ¹	10.1	23.0	32.9	9.86	DNA ²	4.71	DNA*	DNA ³	8.09	4.70
			Max	DNA ¹	23.8	42.2	50.6	25.6	DNA ²	6.96	DNA*	DNA ³	27.3	13.6
			Min	DNA ¹	3.12	11.8	14.2	1.73	DNA ²	3.17	DNA*	DNA ³	1.53	0.86
			Excedance(Days)	DNA ¹	0	0	0	0	DNA ²	0	DNA*	DNA ³	0	0
			Data capture(%)	DNA ¹	82	98	97	86	DNA ²	99	DNA*	DNA ³	76	97
NO ₂ -24 hr	ppb	53 (Annual)	Average	DNA ¹	DNA*	66.0	DNA*	29.0	DNA ²	33.4	35.9	DNA ³	DNA*	18.3
			Max	DNA ¹	DNA*	136	DNA*	51.3	DNA ²	101	63.2	DNA ³	DNA*	35.3
			Min	DNA ¹	DNA*	16.7	DNA*	4.10	DNA ²	10.7	20.2	DNA ³	DNA*	13.2
			Excedance(Days)	DNA ¹	DNA*	17	DNA*	0	DNA ²	3	2	DNA ³	DNA*	0
			Data capture(%)	DNA ¹	DNA*	96	DNA*	96	DNA ²	97	100	DNA ³	DNA*	98
CO- 1 hr	ppm	35	Average	DNA ¹	2.43	4.01	DNA*	DNA*	DNA ²	2.37	3.91	DNA ³	DNA*	DNA*
			Max	DNA ¹	13.3	11.3	DNA*	DNA*	DNA ²	7.12	9.64	DNA ³	DNA*	DNA*
			Min	DNA ¹	0.05	2.36	DNA*	DNA*	DNA ²	0.14	2.50	DNA ³	DNA*	DNA*
			Excedance(Hour)	DNA ¹	0	0	DNA*	DNA*	DNA ²	0	0	DNA ³	DNA*	DNA*
			Data capture(%)	DNA ¹	97	98	DNA*	DNA*	DNA ²	99	100	DNA ³	DNA*	DNA*
CO-8hr	ppm	9	Average	DNA ¹	2.42	4.00	DNA*	DNA*	DNA ²	2.36	3.90	DNA ³	DNA*	DNA*
			Max	DNA ¹	12.1	6.78	DNA*	DNA*	DNA ²	5.16	6.97	DNA ³	DNA*	DNA*
			Min	DNA ¹	0.29	2.47	DNA*	DNA*	DNA ²	1.03	2.60	DNA ³	DNA*	DNA*
			Excedance(Hour)	DNA ¹	19	0	DNA*	DNA*	DNA ²	0	0	DNA ³	DNA*	DNA*
			Data capture(%)	DNA ¹	98	97	DNA*	DNA*	DNA ²	97	99	DNA ³	DNA*	DNA*
O ₃ -1hr	ppb	120	Average	DNA ¹	DNA*	5.09	DNA*	DNA*	DNA ²	15.6	10.1	DNA ³	13.7	DNA*
			Max	DNA ¹	DNA*	28.3	DNA*	10.6	DNA ²	66.4	51.7	DNA ³	53.7	DNA*
			Min	DNA ¹	DNA*	1.29	DNA*	1.08	DNA ²	0.53	0.05	DNA ³	0.96	DNA*
			Excedance(Hour)	DNA ¹	DNA*	0	DNA*	0	DNA ²	0	0	DNA ³	0	DNA*
			Data capture(%)	DNA ¹	DNA*	97	DNA*	96	DNA ²	98	69	DNA ³	99	DNA*
O ₃ -8hr	ppb	80	Average	DNA ¹	DNA*	5.12	DNA*	2.44	DNA ²	15.7	10.1	DNA ³	13.8	DNA*
			Max	DNA ¹	DNA*	20.2	DNA*	6.68	DNA ²	53.7	41.3	DNA ³	43.7	DNA*
			Min	DNA ¹	DNA*	1.41	DNA*	1.22	DNA ²	1.67	0.08	DNA ³	1.58	DNA*
			Excedance(Hour)	DNA ¹	DNA*	0	DNA*	0	DNA ²	0	0	DNA ³	0	DNA*
			Data capture(%)	DNA ¹	DNA*	95	DNA*	94	DNA ²	97	68	DNA ³	98	DNA*

CAMS= Continuous Air Monitoring Station, NAAQS=National Ambient Air Quality Standard, a=Refurbishment CAMS, PM= Particulate Matter

DNA= Data Not Available, 1-3= DNA due to station not within monitoring network due to malfunction of DAS, *=DNA due to malfunction of the analyzer/sensor/ poor data capture rate

Table 4: Summary Air Quality and Meteorological data measured during December, 2016 at different CAMS operated under DoE (Cont'd)

Parameter	unit	NAAQS	Summary	CAMS-1 (S-Bhaban)	CAMS-2 (BARC) ^a	CAMS-3 (D-salam)	CAMS-4 (Gazipur)	CAMS-5 (Narayong anj)	CAMS-6 TV-St (Chittagong) ^a	CAMS-7 Agrabad- (Chittagong)	CAMS-8 (Sylhet)	CAMS-9 (Khulna) ^a	CAMS-10 (Rajshahi) ^a	CAMS-11 (Barisal)
PM _{2.5} -24hr	µg /m ³	65	Average	DNA ¹	DNA*	169	161	206	DNA ²	139	86.5	DNA ³	121	141
			Max	DNA ¹	DNA*	269	271	319	DNA ²	280	170	DNA ³	306	229
			Min	DNA ¹	DNA*	85.9	82.3	122	DNA ²	69.5	54.9	DNA ³	46.7	80.3
			Excedance(Days)	DNA ¹	DNA*	21	30	23	DNA ²	29	11	DNA ³	23	29
			Data capture(%)	DNA ¹	DNA*	70	93	74	DNA ²	96	73	DNA ³	90	97
PM ₁₀ -24hr	µg /m ³	150	Average	DNA ¹	DNA*	272	238	345	DNA ²	205	160	DNA ³	183	177
			Max	DNA ¹	DNA*	447	394	552	DNA ²	345	277	DNA ³	277	213
			Min	DNA ¹	DNA*	163	173	143	DNA ²	112	95	DNA ³	108	129
			Excedance(Days)	DNA ¹	DNA*	21	28	29	DNA ²	23	14	DNA ³	24	13
			Data capture(%)	DNA ¹	DNA*	69	91	91	DNA ²	92	98	DNA ³	98	47
Solar rad. 1hr	watt/m ²	NA	Average	DNA ¹	DNA*	134	144	DNA*	DNA ²	139	162	DNA ³	DNA*	135
			Max	DNA ¹	DNA*	677	709	DNA*	DNA ²	619	692	DNA ³	DNA*	654
			Min	DNA ¹	DNA*	7.55	5.74	DNA*	DNA ²	7.00	6.85	DNA ³	DNA*	8.13
			Data capture(%)	DNA ¹	DNA*	99	96	DNA*	DNA ²	99	100	DNA ³	DNA*	98
Relative Humidity 1hr	(%)	NA	Average	DNA ¹	DNA*	71.1	77.3	68.1	DNA ²	70.0	71.5	DNA ³	DNA*	79.8
			Max	DNA ¹	DNA*	94.3	96.8	91.7	DNA ²	93.3	99.4	DNA ³	DNA*	99.5
			Min	DNA ¹	DNA*	30.4	38.8	28.6	DNA ²	33.3	29.6	DNA ³	DNA*	32.8
			Data capture(%)	DNA ¹	DNA*	98	96	96	DNA ²	99	100	DNA ³	DNA*	98
Ambient Temp. 1hr	(°c)	NA	Average	DNA ¹	28.5	22.0	21.3	DNA*	DNA ²	22.9	22.2	DNA ³	DNA*	22.6
			Max	DNA ¹	36.4	30.8	44.9	DNA*	DNA ²	38.0	30.1	DNA ³	DNA*	33.3
			Min	DNA ¹	25.8	15.2	14.5	DNA*	DNA ²	16.6	16.3	DNA ³	DNA*	15.2
			Data capture(%)	DNA ¹	99	98	98	DNA*	DNA ²	99	100	DNA ³	DNA*	98
Rainfall 1hr	(m.m.)	NA	Average	DNA ¹	0.72	0.02	DNA*	0.25	DNA ²	0.02	DNA*	DNA ³	DNA*	DNA*
			Max	DNA ¹	2.61	0.05	DNA*	0.60	DNA ²	0.03	DNA*	DNA ³	DNA*	DNA*
			Min	DNA ¹	0.03	0.02	DNA*	0.10	DNA ²	0.02	DNA*	DNA ³	DNA*	DNA*
			Data capture(%)	DNA ¹	64	32	DNA*	90	DNA ²	57	DNA*	DNA ³	DNA*	DNA*

CAMS= Continuous Air Monitoring Station, NAAQS=National Ambient Air Quality Standard, a=Refurbishment CAMS, PM= Particulate Matter

DNA= Data Not Available, 1-3= DNA due to station not within monitoring network due to malfunction of DAS, *=DNA due to malfunction of the analyzer/sensor/ poor data capture rate

FIGURE 3: TIME SERIES OF ALL PARAMETERS (SO₂, NO_x AND O₃) MEASURED IN ALL CAMS DURING DECEMBER, 2016







Figure 4: TIME SERIES OF ALL PARAMETERS (CO, PM10 AND PM2.5) MEASURED IN CAMS DURING DECEMBER, 2016







Figure 5: Monthly Summary of AQI for month of December, 2016

